

🔥 **Pack Size** - 20, 1000 litres

Analysis	w/w	w/v
Nitrogen (N)	10.0%	14.0%
Phosphate (P <sub>2</sub> O <sub>5</sub> )	33.5%	46.0%
Zinc	0.18%	0.25%

🔥 **Directions for Use**

Use Folex PZ when phosphate deficiency is diagnosed or suspected, when crop demand is high and cannot be met by soil reserves or as part of a nutrient management programme. Foliar uptake will be enhanced by the addition of NA13<sup>1</sup> unless already in tank mix with a pesticide.

Apply 7-15 l/ha, in a minimum of 200 l/ha water.

The spray tank should be filled with half the required water. If applicable, add the required amount of NA13 to the water before the Folex PZ. Measure the required amount of Folex PZ and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate l/ha	Comments
Maize	4-8 leaves	7	Apply when deficiency is suspected, when soil/weather conditions prevent adequate phosphate uptake through the roots, or when SAP analysis shows low nutrient status  Repeat as necessary at 10-14 day intervals
Potatoes	7-10 days after tuber initiation	15	At 7-10 days after start of tuber initiation crops are usually meeting along the rows at this stage

🔥 **Notes**

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility, tank mixing and for physical compatibility with pesticides refer to the website [www.omex.co.uk](http://www.omex.co.uk)

<sup>1</sup> NA13 is an adjuvant designed to help with improved adhesion, deposition and penetration of the spray solution on the leaf surface. NA13 should be added at 0.1% of the spray volume, e.g. 100 ml in 100 litres of water. Maintain agitation and apply immediately after mixing. See website [www.omex.co.uk](http://www.omex.co.uk) for details.