

💧 **Pack Size** - 10,1000 litres

Analysis	w/w	w/v
Copper	5.0%	6.0%
Sulphur (SO ₃)	6.5%	7.5%

💧 **Directions for Use**

Use Folex Cu 6 when deficiency is diagnosed or suspected on soils where conditions are likely to cause copper deficiency in the crop. Foliar uptake will be enhanced by the addition of NA13¹ unless already in tank mix with a pesticide.

Apply 1-2 L/ha, in a minimum of 200 L/ha water.

The spray tank should be filled with half the required water. If applicable, add the required amount of NA13 to the water before the Folex Cu 6. After shaking the container, measure the required amount of Folex Cu 6 and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate L/ha	Comments
Cereals	Spring	1	Apply where known deficiency exists before symptoms appear
	Late spring and summer	2	Apply when symptoms appear and repeat as necessary
Onions, Leeks	Once crop is established	2	A second application may be required in dry seasons
Other crops	As required	2	Apply where known deficiency exists before symptoms appear
		2	Repeat as necessary

💧 **Notes**

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility and tank mixing refer to the section on pages 86-87, and for physical compatibility with pesticides refer to the website www.omex.co.uk

¹ NA13 is an adjuvant designed to help with improved adhesion, deposition and penetration of the spray solution on the leaf surface. NA13 should be added at 0.1% of the spray volume, e.g. 100 ml in 100 litres of water. Maintain agitation and apply immediately after mixing. See page 82 for details.