

💧 Pack Size - 10,1000 litres

Analysis	w/w	w/v
Sulphur (SO ₃)	14.0%	18.5%
Zinc	11.0%	15.0%

💧 Directions for Use

Use Folex Zn when zinc deficiency is diagnosed or suspected. Foliar uptake will be enhanced by the addition of NA13¹ unless already in tank mix with a pesticide.

Apply 1-2 L/ha, in a minimum of 200 L/ha water.

The spray tank should be filled with half the required water. If applicable, add the required amount of NA13 to the water before the Folex Zn. Measure the required amount of Folex Zn and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate L/ha	Comments
Brassicas	4 - 6 leaves	1 - 2	Repeat as necessary
Maize	4 - 8 leaves	1 - 2	Apply early to aid rapid growth Repeat 14-21 days later
Onions & Leeks	3 - 4 true leaves	1 - 2	Apply early to aid rapid growth Repeat 14-21 days later
Potatoes	20cm high	1 - 2	Repeat as necessary

💧 Notes

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility and tank mixing refer to the section on pages 86-87, and for physical compatibility with pesticides refer to the website www.omex.co.uk

¹ NA13 is an adjuvant designed to help with improved adhesion, deposition and penetration of the spray solution on the leaf surface. NA13 should be added at 0.1% of the spray volume, e.g. 100 ml in 100 litres of water. Maintain agitation and apply immediately after mixing. See page 82 for details.