

♦ Pack Size - 10,1000 litres

Analysis	w/w	w/v
Sulphur (S)	51.0%	72.0%
Sulphur (SO ₃)	128.0%	180.0%

♦ Directions for Use: Foliar Applications

Apply 5-10 L/ha, in a minimum of 200 L/ha water.

The spray tank should be filled with half the required water. After shaking the container, measure amount of Sulflo and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate L/ha	Comments
Cereals	Autumn/Spring	10	Apply in autumn if sufficient crop cover
Oilseed Rape	Start of stem extension	5	Maintenance, apply early during maximum growth
		10	Deficiency
Grass	Autumn/Spring	10	After 1st cut and 2nd cut
Potatoes	During Bulking	5	Deficiency
Sugar Beet	End of July	5-10	Helps reduce amino-N levels

♦ Directions for Use: Pre-Planting/In Furrow Applications

To reduce soil pH or to improve skin finish apply 40-100 L/ha, in 200-400 L/ha water/liquid fertiliser.

Crop	Timing	Rate L/ha	Comments
Potatoes	Immediately before or during planting	40 - 100	Apply post ploughing but pre-planting to achieve an even distribution through the ridge. Use the higher rate on high pH soils

♦ Notes

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility and tank mixing refer to the section on pages 86-87, and for physical compatibility with pesticides refer to the website www.omex.co.uk