

Folex® Range

Foliar Nutrition - Boron

Use

To correct micronutrient deficiency

Crops

Most horticultural crops.

Pack Size

10 litres

Function of Folex Range

A range of single micronutrient supplements formulated as sulphates or in chelated form..



Nutrient	Product	Analysis w/w	Analysis w/v
Boron (B)	Folex B	11.0%	15.0%
Copper Sulphate (Cu)	Folex Cu 6	5.0%	6.0%
Magnesium Sulphate (MgO)	Folex Mg 9	7.3%	9.0%
Manganese Sulphate (Mn)	Manganese 17.5	12.3%	17.5%
Zinc Sulphate (Zn)	Folex Zn	11.0%	15.0%

🔥 Directions for Use

Applied as directed below. Foliar uptake will be enhanced by the addition of NA13¹ unless a spray tank mix with a pesticide.



Apply 200-600 L/ha water.

The spray tank should be filled with half the required water. If applicable, add the required amount of NA13 to the water before the nutrient. After shaking the container, measure the required amount of product and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate L/ha				
		Folex B	Folex Cu 6	Folex Mg 9	Mn 17.5	Folex Zn
Protected Edibles	When deficiency is diagnosed or suspected, or prior to rapid growth	2	2	2.5	2	1
Field Vegetables	When deficiency is diagnosed or suspected, or prior to rapid growth	2	2	7.5	3	1
Bulbs & Outdoor Flowers	From 10cm high but before flowering	2	2	7.5	3	1
Hardy Nursery Stock	Apply to new season growth or when deficiency is diagnosed or suspected	N/A	2	3	2	1
Protected Ornamentals	When deficiency is diagnosed or suspected, or prior to rapid growth	2	2	2	2	1
Soft Fruit	From start of flowering to fruit set	2	1	3	3	1
Tree Fruit	From petal fall (copper - after harvest, manganese - before flowering)	2	0.5	7.5	3	2

🔥 Notes

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility and tank mixing refer to the section on pages 66-67, and for physical compatibility with pesticides refer to the website www.omex.co.uk

¹ NA13 is an adjuvant designed to help with improved adhesion, deposition and penetration of the spray solution on the leaf surface. NA13 should be added at 0.1% of the spray volume, e.g. 100 ml in 100 litres of water. Maintain agitation and apply immediately after mixing.

For more information contact: OMEX Horticulture, Estuary Road, King's Lynn, Norfolk, PE30 2HH

Web www.omex.co.uk • **Email** horticulture@omex.com • **Tel** 01553 760011