

Sulphomex®

Foliar Nutrition - Ammonium Thiosulphate

Use

To correct nutrient deficiency.

Crops

Most field grown horticultural crops.

Pack Size

20, 1000 litres



Analysis	w/w	w/v
Nitrogen (N)	11.3%	15.0%
Sulphur (SO ₃)	65.0%	87.5%

◆ Directions for Use

Use Sulphomex to correct sulphur deficiency, or as part of a programme to prevent sulphur deficiency.

Apply 5-10 L/ha, in a minimum of 200 L/ha water.

The spray tank should be filled with half the required water. After shaking the container, measure the required amount of Sulphomex and add to the tank whilst maintaining constant agitation. Add remaining water to correct dilution and spray.

Crop	Timing	Rate L/ha	Comments
Brassicas	From 4-6 true leaves	5	Maintenance, apply early during maximum growth
		10	
Blueberries (dormant)	Apply when the crop is dormant	3	Apply post pruning, but before blossom and at a minimum rate of 1L/200L/ha water and a maximum rate of 3L/ha in a minimum of 600L water/ha.
Other Field Vegetables	When crop is 15cm tall, or 6 true leaves	5	Repeat after 10-14 days if deficiency is severe

◆ Notes

Do not apply in tank mix with pesticides when crop is showing deficiency symptoms, is under stress, or in adverse weather conditions.

For further information on compatibility, tank mixing refer and physical compatibility with pesticides refer to the website www.omex.com